

LAKE MACQUARIE – WYONG REVIEW OF MONTHLY AMBIENT AIR QUALITY DATA JUNE 2015

NSW Environment Protection Authority

29 September 2015

Job Number 14030303A

Prepared by

Todoroski Air Sciences Pty Ltd

Suite 2B, 14 Glen Street Eastwood, NSW 2122 Phone: (02) 9874 2123

Email: info@airsciences.com.au

(02) 9874 2125



Lake Macquarie – Wyong Review of Monthly Ambient Air Quality

Data

June 2015

Author(s): Aleks Todoroski Dan Kjellberg

Position: Director Atmospheric Scientist

DOCUMENT CONTROL

Report Version	Date	Prepared by	Reviewed by	
DRAFT - 001	29/09/2015	D. Kjellberg		
FINAL - 001	29/09/2015	D. Kjellberg	A. Todoroski	

This report has been prepared in accordance with the scope of works between Todoroski Air Sciences Pty Ltd (TAS) and the client. TAS relies on and presumes accurate the information (or lack thereof) made available to it to conduct the work. If this is not the case, the findings of the report June change. TAS has applied the usual care and diligence of the profession prevailing at the time of preparing this report and commensurate with the information available. No other warranty or guarantee is implied in regard to the content and findings of the report. The report has been prepared exclusively for the use of the client, for the stated purpose and must be read in full. No responsibility is accepted for the use of the report or part thereof in any other context or by any third party.

© Copyright State of NSW and the Environment Protection Authority



TABLE OF CONTENTS

1	INTF	RODUCTION	1
2		JECT SCOPE	
3	THE	PURPOSE OF AMBIENT MONITORING	1
	3.1	More about air quality	2
4	AIR	QUALITY MONITORING SITES	3
5	AIR	QUALITY CRITERIA	4
	5.1	Particulate matter	4
	5.1.1	PM _{2.5} concentrations	4
	5.2	Other air pollutants	4
	5.3	Summary of applicable criteria for this assessment	5
6	MET	EOROLOGICAL MONITORING DATA	5
7	AME	BIENT AIR QUALITY MONITORING DATA	7
	7.1	Preamble	7
	7.2	Analysis of Monitoring Data	7
	7.3	PM ₁₀	7
	7.4	PM _{2.5}	8
	7.5	Nitrogen dioxide NO ₂	8
	7.6	Sulfur dioxide SO ₂	8
8	ANA	LYSIS OF ELEVATED POLLUTANT LEVELS	.13
9	CON	ICLUSIONS	.14
1() REFE	ERENCES	.15



LIST OF TABLES

Table 4-1: Monitoring sites	3
Table 5-1: EPA air quality impact assessment criteria	
Table 5-2: Advisory standard for PM _{2.5} concentrations	4
Table 5-3: Air quality impact assessment criteria for air pollutants	5
Table 5-4: Air quality impact assessment criteria used in this assessment	5
Table 7-1: Maximum pollutant levels - June 2015	7
LIST OF FIGURES	
Figure 4-1: Monitoring site locations	3
Figure 6-1: June windroses – Wallsend, Dora Creek, Marks Point, Wyee, Norah Head and Wyong	6
Figure 7-1: Lake Macquarie - Wyong 24-hour average PM ₁₀ levels - June 2015	
Figure 7-2: Lake Macquarie - Wyong 24-hour average PM _{2.5} levels – June 2015	10
Figure 7-3: Lake Macquarie - Wyong 1-hour average NO ₂ levels - June 2015	11
Figure 7-4: Lake Macquarie - Wyong 1-hour average SO ₂ levels – June 2015	12

LIST OF APPENDICIES

Appendix A – How to read a windrose


Appendix B – Monitoring Data (Graphical)

Appendix C – Monitoring Data (Tabulated)

EXECUTIVE SUMMARY

This report has been prepared by Todoroski Air Sciences for the NSW Environment Protection Authority (NSW EPA) and presents ambient air quality monitoring data recorded in the Lake Macquarie - Wyong region for the month of June 2015. The results indicate that the air quality was generally very good in the Lake Macquarie - Wyong region during June.

The data summary (shown below) indicates that in June 2015, all data were below the applicable criteria. Further details are provided in the report. The 24-hour average data are provided in the Appendices.



Lake Macquarie - Wyong Air Quality Pictorial Summary - June 2015

Lake Macquarie - Wyong Air Quality Tabular Summary - June 2015

	PM ₁₀ (μg/m³)	PM _{2.5} (μg/m ³)	SO₂ (μg/m³)	NO ₂ (μg/m³)	SO₂ (μg/m³)			
Site	24-hour average	24-hour average	24-hour average	1-hour average	1-hour average			
Site	Air Quality Impact Criteria							
	50	25*	228	246	570			
Wallsend	✓	✓	✓	✓	✓			
Wyong	✓	✓	✓	✓	✓			
Dora Creek	-	-	✓	✓	✓			
Marks Point	-	-	✓	✓	✓			
Wyee	-	✓	✓	✓	✓			
Wakefield HVAS	✓	-	-	-	-			
Teralba HVAS	✓	-	-	-	-			

^{✓ -} All data below applicable criteria

x - At least one elevated level above applicable criteria

⁻ Not applicable
HVAS - High Volume Air Sampler

Advisory reporting standard for PM_{2.5} concentrations (refer to Section 5.1)

1 INTRODUCTION

This report has been prepared by Todoroski Air Sciences on behalf of the NSW EPA. It provides a summary and analysis of the available ambient air quality and meteorological data collected in the Lake Macquarie - Wyong region during June 2015.

2 PROJECT SCOPE

The following outlines the scope of work for this project.

- Provide a monthly report written in plain English to the NSW EPA summarising and analysing available air quality data and meteorological information.
- + The report will be published on the EPA's website and will assess the available data from monitoring stations operated by the NSW Office of Environment and Heritage (OEH) at Wyong and Wallsend, and by industry at Wyee, Marks Point, Dora Creek, Wakefield and Teralba.
- → The aim is to provide a simplified report that is accessible and contains results that would be clearly understood by the general public.

The work is for the period from September 2013 to June 2015.

3 THE PURPOSE OF AMBIENT MONITORING

It is important to note that the data presented in this report are from both EPA and Industry monitoring sites. The EPA and the industry sites collect data for different purposes and this needs to be understood when comparing the data to the criteria.

EPA monitoring sites are specifically designed to measure the likely levels of pollutants that the general population in the area would experience (i.e. an underlying population exposure level), whereas industry monitoring sites are specifically designed to measure maximum levels in a particular location that May be affected by a particular industry.

Data from EPA sites can be compared with national air quality standards. Where the levels measured at EPA monitoring sites are above the national standards on a prolonged and consistent basis, this indicates that some investigation of the potential cause of the issue may be warranted to determine whether any action on a regional level would reduce or better manage the pollutant levels. In the case of PM₁₀, it is noted that the national standards permit five days annually above the criteria to allow for events such as bushfires and dust storms.

Data from industry monitoring sites can be compared with EPA impact assessment criteria. Where the levels measured at industry monitoring sites are above the impact assessment criteria on a prolonged and consistent basis, this indicates that further investigation is warranted to determine whether industry is responsible, and if so whether action to reduce or better manage the pollutant can be taken.

Whether there is any harmful effect on an individual due to an air pollutant will depend on many additional factors, and not just on the measured level of a pollutant. These factors include the total exposure to the pollutant, individual circumstances (age, health, body mass, levels of pollutants at work), levels of other pollutants in the area, and many other factors.

Where pollutant levels are below the criteria generally, harm would not be expected to occur, but it does not follow that harm automatically occurs when pollutant levels are above the criteria.

The criteria serve to highlight potential issues with the levels of pollutants that may warrant more detailed examination. The criteria may also serve to prioritise action in various areas, for example areas with the highest pollutant levels and highest populations or highest exposure would be expected to receive priority action.

3.1 More about air quality

More information about air quality can be found via the following links:

- + The Air Quality Index (AQI) was developed by the NSW EPA as an easily understood means of rating the pollutant level relative to its pollutant criteria.
 - o http://www.environment.nsw.gov.au/AQMS/aboutaqi.htm
- + Aqicn.org provides a near real-time AQI values for monitoring locations around the world. It should be noted that the AQI presented on this website is calculated differently to the NSW EPA AQI and is less stringent than those used in Australia, thus a direct comparison may not be valid.
 - http://agicn.org/map/world/
- + The NSW OEH website air quality page provides hourly updates of the AQI and data readings from the NSW EPA monitoring sites, and can provide daily forecasts for Sydney and alerts for elevated levels at Wallsend a Wyong, for example. The web tool also presents near real-time wind and pollutant data readings overlaid on regional maps for the Upper Hunter and Newcastle.
 - o http://www.environment.nsw.gov.au/aqms/aqi.htm
- + The Lower Hunter Particle Characterisation Study aims to determine the composition of particulate samples collected at monitoring sites at Beresfield, Newcastle, Stockton and Mayfield, and to identify the potential major sources of fine particulates in Newcastle and the Lower Hunter. Progress reports are published on the OEH website provided below.
 - http://www.environment.nsw.gov.au/aqms/lowhunterparticle.htm
- + The Air Emissions in My Community web tool presents the estimated emission quantities of various substances and their sources by postcode (and larger) sized areas in an easy to use graphical interface. This is one of the best inventories of emissions that is available, but it is important to appreciate that it cannot include all sources of emissions. It is important to also understand that pollutant emissions are not the same as the pollutant levels that this report presents. Emissions in a given area are one of several important factors that affect pollutant levels in an area, for example the dispersion of the emissions in the atmosphere and how the emissions are released are critical in determining the air quality pollutant levels.
 - o http://www.epa.nsw.gov.au/air/airemissionsapp/airemissionswebtool.aspx
- + The NSW Health website provides information on how air pollution affects health and steps for reducing your air pollution and limiting your exposure.
 - o http://www.health.nsw.gov.au/environment/air/Pages/default.aspx

4 AIR QUALITY MONITORING SITES

Figure 4-1 and **Table 4-1** summarise the locations and recorded parameters of the monitoring sites in the Lake Macquarie - Wyong region in June 2015.




Figure 4-1: Monitoring site locations

Table 4-1: Monitoring sites

Table 4-1: Monitoring sites								
Monitoring Station	Туре	Recorded Parameters	Recording Periods					
Wallsend	NSW EPA site	PM ₁₀ (TEOM), PM _{2.5} , NO ₂ , SO ₂ , WS, WD	Hourly/Daily					
Wyong	NSW EPA site	PM ₁₀ (TEOM), PM _{2.5} , NO ₂ , SO ₂ , WS, WD	Hourly/Daily					
Marks Point	Industry site	NO ₂ , SO ₂ , WS, WD	Hourly					
Wyee	Industry site	PM _{2.5} , NO ₂ , SO ₂ , WS, WD	Hourly					
Dora Creek	Industry site	NO ₂ , SO ₂ , WS, WD	Hourly					
Norah Head	BOM weather station	WS, WD	Hourly					
Wakefield HVAS	Industry site	PM ₁₀ (HVAS)	Every 6th Day					
Teralba HVAS	Industry site	PM ₁₀ (HVAS)	Every 6th Day					

 PM_{10} - Particulate matter < $10\mu m$

 $PM_{2.5}$ - Particulate matter < 2.5 μ m

TEOM - Tapered Element Oscillating Microbalance (which samples air continuously)

NO₂ - Nitrogen dioxide

SO₂ - Sulfur dioxide

HVAS - High volume air sampler (which samples for a 24-hour period every 6 days)

WS - Wind speed WD - Wind direction BOM - Bureau of Meteorology

5 AIR QUALITY CRITERIA

The sections below identify the key pollutants currently being monitored at the Lake Macquarie - Wyong air quality monitoring sites and the applicable air quality criteria.

5.1 Particulate matter

Particulate matter consists of particles of varying size and composition. The total mass of all particles suspended in air is defined as the Total Suspended Particulate matter (TSP). The upper size range for TSP is nominally taken to be 30 micrometres (μ m) as in practice particles larger than 30 to 50 μ m will settle out of the atmosphere too quickly to be regarded as air pollutants.

The TSP is defined further into two sub-components. They are PM_{10} particles, particulate matter with aerodynamic diameters of $10\mu m$ or less, and $PM_{2.5}$, particulate matter with aerodynamic diameters of $2.5\mu m$ or less.

Table 5-1 summarises the air quality goals that are relevant to particulate pollutants as outlined in the NSW Environment Protection Agency (EPA) document "Approved Methods for the Modelling and Assessment of Air Pollutants in NSW" (NSW DEC, 2005).

Table 5-1: EPA air quality impact assessment criteria

Pollutant	Averaging Period	Criterion
Total suspended particulates (TSP)	Annual	90μg/m³
Douticulate Matter (100m (DM)	Annual	30μg/m³
Particulate Matter $< 10\mu m (PM_{10})$	24-hour	50μg/m³

Source: NSW DEC, 2005

5.1.1 PM_{2.5} concentrations

The NSW EPA currently do not have impact assessment criteria for PM_{2.5} concentrations, however the National Environment Protection Council (NEPC) has released a variation to the National Environment Protection Measure (NEPM) (**NEPC**, **2003**) to include advisory reporting standards for PM_{2.5} (see **Table 5-2**). As with the NEPM goals, the advisory reporting standards apply to the average, or general exposure of a population, rather than to "hot spot" locations such as industry monitoring sites.

Table 5-2: Advisory standard for PM_{2.5} concentrations

Pollutant	Averaging Period	Concentration
Particulate Matter < 2 Fum (PM.)	24-hour	25μg/m³
Particulate Matter < $2.5\mu m$ (PM _{2.5})	Annual	8µg/m³

Source: NEPC, 2003

5.2 Other air pollutants

Nitrogen dioxide (NO_2) is reddish-brown in colour (at high concentrations) with a characteristic odour and can irritate the lungs and lower resistance to respiratory infections such as influenza. NO_2 belongs to a family of reactive gases called nitrogen oxides (NO_x). These gases form when fuel is burned at high temperatures, and mainly originates from motor vehicles, power generators and industrial boilers (**USEPA, 2013**). NO_x may also be generated by blasting activities. It is important to note that when formed, NO_2 is generally a small fraction of the total NO_x generated.

Sulfur dioxide (SO₂) is a colourless, toxic gas with a pungent and irritating smell. It commonly arises in industrial emissions due to the sulfur content of the fuel. SO₂ can have impacts upon human health

and the habitability of the environment for flora and fauna. SO_2 emissions are a precursor to acid rain, which can be an issue in the northern hemisphere; however it is not known to be an issue in NSW.

Table 5-3 summarises the air quality goals for NO₂ and SO₂.

Table 5-3: Air quality impact assessment criteria for air pollutants

Pollutant	Averaging period	Criterion
Nitrogen Dioxide (NO ₂)	1-hour	246μg/m³
Nitrogen Dioxide (NO ₂)	Annual	62μg/m³
	10-minute	712μg/m³
Sulfur Dioxide (SO ₂)	1-hour	570μg/m³
Sulful Dioxide (SO ₂)	24-hour	228μg/m³
	Annual	60μg/m³

Source: NSW DEC, 2005

5.3 Summary of applicable criteria for this assessment

The particulate and gaseous pollutants monitored in the Lake Macquarie – Wyong region have air quality criteria which are averaged over short and long time periods. Annually averaged criteria require a full year of data.

As this report only looks at one month of ambient air quality data, the annual average criteria are not applicable. The SO_2 10-minute average criterion was not included as 10-minute monitoring data are not available. Therefore the criteria relevant to this assessment are those averaged over the shorter time periods (1-hour and 24-hours).

Table 5-4 summarises the applicable air quality criteria for this assessment.

Table 5-4: Air quality impact assessment criteria used in this assessment

Pollutant	Averaging Period	Туре	Concentration	
Particulate Matter $< 10 \mu m$ (PM ₁₀)	24-hour	Criterion	50μg/m³	
Particulate Matter < 2.5μm (PM _{2.5})	24-hour	Advisory Reporting Standard	25μg/m³	
Nitrogen Dioxide (NO ₂)	1-hour	Criterion	246μg/m³	
Sulfur Dioxide (SO ₂)	1-hour	Criterion	570μg/m³	
Sullul Dioxide (SO ₂)	24-hour	Criterion	228μg/m³	

6 METEOROLOGICAL MONITORING DATA

Representative wind speed and direction data have been obtained from the Lake Macquarie - Wyong air quality monitoring stations. The data are presented as a series windroses. For an example of how to read a windrose, refer to **Figure A-1** in **Appendix A**.

Figure 6-1 presents the June 2015 windroses for Wallsend, Dora Creek, Marks Point, Wyee, Norah Head and Wyong.

The figure shows that the meteorological stations recorded winds which varied depending on the local influence of environmental features such as terrain, vegetation and buildings. Overall the stations recorded low wind speeds which typically originated from the north-westerly and south-westerly quadrants.

The Norah Head weather station recorded wind speeds which were generally higher than those recorded at the other stations. This is expected as the Norah Head weather station is located in an unsheltered coastal location that would be largely influenced by sea breezes.




Figure 6-1: June windroses – Wallsend, Dora Creek, Marks Point, Wyee, Norah Head and Wyong

The meteorological stations generally recorded low wind speeds which originated from the north-westerly and south-westerly quadrants.

AMBIENT AIR QUALITY MONITORING DATA

7.1 Preamble

The monitoring data in this report are presented in raw form as provided to Todoroski Air Sciences by the NSW EPA.

The 24-hour average data presented in this report have been averaged using the 1-hour average readings. Days which contain less than 75% data (less than 18 hours of 1-hour average data) have not been included in this report.

All of the monitoring data provided to Todoroski Air Sciences are presented in this report. The data are shown in the results and appendices as relevant. Hourly data are presented in a graphical format in Appendix B and 24-hour average data are presented in tabulated format in Appendix C.

On 5 June 2015 the Wyee PM_{2.5} monitor underwent annual calibration and maintenance. This included the replacement of all consumable items and the installation of a new pump.

7.2 Analysis of Monitoring Data

Table 7-1 presents a summary of the maximum pollutant levels measured during June 2015. The results indicate that pollutant levels were below the applicable criteria for all monitors at all times.

PM₁₀ NO₂ SO₂ PM_{2.5} SO₂ $(\mu g/m^3)$ $(\mu g/m^3)$ $(\mu g/m^3)$ $(\mu g/m^3)$ $(\mu g/m^3)$ Site 24-hour 24-hour 24-hour 1-hour 1-hour average average average average average **Air Quality Impact Criteria** 25* 246 50 228 570 Wallsend 22.5 22.0 11.6 49.1 78.3 Wyong 15.2 7.8 6.2 57.0 39.6 Dora Creek 3.9 164.0 7.4 Marks Point 8.9 53.8 55.1 13.2 7.7 35.6 Wvee 30.3 Wakefield HVAS 13.9 Teralba HVAS 20.0

Table 7-1: Maximum pollutant levels - June 2015

7.3 PM₁₀

Figure 7-1 presents all of the 24-hour average PM₁₀ monitoring results recorded in the Lake Macquarie - Wyong region in June 2015.

Relative to the Air Quality Index, as shown by the coloured bands in the figure, PM₁₀ levels were very good at all monitors in June 2015 with the exception of the Wallsend monitor which recorded good levels 37% of the time and the Teralba monitor which recorded good levels on one day.

All data recorded at the Lake Macquarie - Wyong monitoring sites were below the PM₁₀ criterion level in June.

Figure B-1 to Figure B-2 in Appendix B present the 1-hour average PM₁₀ data in graphical form for each individual site. There is no criterion that applies to 1-hour average PM₁₀ levels and these 1-hour

Advisory reporting standard for PM_{2.5} concentrations (refer to Section 5.1)

⁻ Not applicable

results are not intended to be compared with the PM_{10} criterion. It is a normal occurrence, and it is expected that in the normal environment 1-hour average PM_{10} levels will fluctuate more significantly than 24-hour average PM_{10} levels.

7.4 PM_{2.5}

Figure 7-2 presents all of the 24-hour average PM_{2.5} monitoring data recorded in the Lake Macquarie - Wyong region in June 2015.

Relative to the Air Quality Index, as shown by the coloured bands in the figure, the data indicate that PM_{2.5} levels were generally very good to good at all monitors at all times with the exception of the Wallsend monitor which recorded fair levels 13% of the time.

All data recorded at the Lake Macquarie - Wyong monitoring sites were below the 24-hour average $PM_{2.5}$ advisory reporting standard of $25\mu g/m^3$ in June 2015.

On 5 June 2015 the Wyee PM_{2.5} monitor underwent annual calibration and maintenance. The PM_{2.5} levels recorded following the maintenance are consistently higher than the levels recorded in previous months. The Wyee data from 5 June onwards are generally similar to the levels recorded at Wallsend.

Figure B-3 to **Figure B-5** in **Appendix B** present the 1-hour average PM_{2.5} data in graphical form for each individual site. There is no criterion that applies to 1-hour average PM_{2.5} levels and these 1-hour results are not intended to be compared with the PM_{2.5} advisory reporting standard. It is a normal occurrence, and it is expected that in the normal environment 1-hour average PM_{2.5} levels will fluctuate more significantly than 24-hour average PM_{2.5} levels.

7.5 Nitrogen dioxide NO₂

Figure 7-3 presents the 1-hour average NO₂ monitoring data recorded in the Lake Macquarie - Wyong region in June 2015.

Relative to the Air Quality Index, as shown by the coloured bands in the figure, the data indicate the NO₂ levels were very good all monitors at all times with the exception of the Dora Creek monitor which recorded good levels approximately 2% of the time

All data were below the applicable criterion on all days.

7.6 Sulfur dioxide SO₂

Figure 7-4 presents the 1-hour average SO₂ monitoring data recorded in the Lake Macquarie - Wyong region in June 2015.

Relative to the Air Quality Index, as shown by the coloured bands in the figure, the data indicate the SO_2 levels were very good all of the time at all of the monitors.

All data were below the applicable criterion on all days.




Figure 7-1: Lake Macquarie - Wyong 24-hour average PM₁₀ levels - June 2015

The recorded PM₁₀ levels were very good at all monitors in June 2015 with the exception of the Wallsend monitor which recorded good levels 37% of the time and the Teralba monitor which recorded good levels on one day. All data recorded at the Lake Macquarie - Wyong monitoring sites were below the 24-hour average criterion of 50µg/m³.




Figure 7-2: Lake Macquarie - Wyong 24-hour average PM_{2.5} levels – June 2015

The recorded $PM_{2.5}$ levels were generally very good to good at all monitors at all times with the exception of the Wallsend monitor which recorded fair levels 13% of the time. All data recorded at the Lake Macquarie - Wyong monitoring sites were below the 24-hour average $PM_{2.5}$ advisory reporting standard of $25\mu g/m^3$.




Figure 7-3: Lake Macquarie - Wyong 1-hour average NO₂ levels – June 2015

All data recorded at the Lake Macquarie - Wyong monitoring sites were below the 1-hour average NO_2 criterion level of $246\mu g/m^3$ in June 2015. Measured levels of NO_2 were very good at all monitors at all times with the exception of the Dora Creek monitor which recorded good levels approximately 2% of the time.




Figure 7-4: Lake Macquarie - Wyong 1-hour average SO₂ levels - June 2015

All data recorded at the Lake Macquarie - Wyong monitoring sites were below the 1-hour average SO₂ criterion level of 570µg/m³ in June 2015. Measured levels of SO₂ were very good at all monitors at all times.

8 ANALYSIS OF ELEVATED POLLUTANT LEVELS

There were no levels above the assessment criteria in June 2015.

CONCLUSIONS

The results indicate that the monitoring stations recorded very good air quality for the majority of June 2015.

Relative to the Air Quality Index:

- → The measured levels of NO₂ were very good at all monitors at all times with the exception of the Dora Creek monitor which recorded good levels 2% of the time;
- The measured levels of SO₂ were very good at all monitors at all times;
- The measured levels of PM_{2.5} were very good to good at all times for all locations with the exception of the Wallsend monitor which recorded fair levels 13% of the time; and,
- The measured PM₁₀ levels were very good at all locations at all times with the exception of the Wallsend monitor which recorded good levels 37% of the time.

On this basis it can be concluded that the air quality in the Lake Macquarie - Wyong region was generally very good in June 2015.

10 REFERENCES

NEPC (2001)

"National Environment Protection (Ambient Air Quality) Measure Technical Paper No. 5 Data Collection and Handling", National Environment Protection Council, May 2001.

NEPC (2003)

"Variation to the National Environment Protection (Ambient Air Quality) Measure for Particles as PM_{2.5}", National Environment Protection Council, May 2003.

NSW DEC (2005)

"Approved Methods for the Modelling and Assessment of Air Pollutants in NSW", Department of Environment and Conservation (NSW), August 2005.

USEPA (2013)

United States Environmental Protection Health Effects of Pollution, Agency website.http://www.epa.gov/region07/air/quality/health.htm, accessed May 2013.

Appendix A How to read a windrose 14030303A_LMWAQ_June2015_150929.docx




Figure A-1: How to read a windrose






Figure B-1: Wallsend PM₁₀ concentration - June




Figure B-2: Wyong PM₁₀ concentration - June




Figure B-3: Wallsend PM_{2.5} concentration – June




Figure B-4: Wyong PM_{2.5} concentration – June




Figure B-5: Wyee PM_{2.5} concentration - June

Appendix C	
Monitoring Data (Tabulated)	
-	140202024 LMWAQ luna201E 1E0020 da sv

Table C-1: June 24-hour average monitoring data

PM ₁₀ (μg/m³)		PM _{2.5} (µg/m³)			SO ₂ (μg/m³)					
Date	Wallsend	Wyong	Wallsend	Wyong	Wyee	Wallsend	Wyong	Dora Creek	Marks Point	Wyee
1/06/2015	7.6	6.6	4.9	2.3	0.5	1.3	0.0	0.7	6.1	1.9
2/06/2015	10.6	11.4	8.2	3.6	1.1	4.3	0.0	1.2	8.6	5.0
3/06/2015	16.8	9.9	13.7	3.4	2.0	11.6	0.1	-	8.4	3.1
4/06/2015	19.5	8.4	17.2	4.3	3.7	10.8	2.9	-	6.7	6.2
5/06/2015	11.8	10.3	8.6	4.8	8.7	-0.2	0.3	-	4.8	2.1
6/06/2015	17.6	15.2	13.2	6.8	11.2	0.7	0.2	3.9	4.9	3.2
7/06/2015	22.5	10.7	22.0	6.5	13.2	4.9	2.9	1.5	8.9	5.1
8/06/2015	13.1	11.6	9.5	7.6	9.9	2.4	0.5	1.0	4.0	2.9
9/06/2015	15.2	11.4	6.3	5.5	8.8	2.3	1.1	1.4	4.3	3.5
10/06/2015	13.1	11.6	5.5	6.6	7.0	0.8	0.5	-	3.5	1.8
11/06/2015	10.2	10.9	5.5	4.4	8.2	1.0	0.5	0.8	4.4	2.5
12/06/2015	13.1	-	7.6	-	9.5	2.0	-	1.6	5.7	5.7
13/06/2015	14.7	6.8	10.1	2.7	9.4	1.3	0.9	2.2	4.2	3.7
14/06/2015	18.1	10.7	16.2	7.4	12.7	2.8	3.2	-	6.5	7.6
15/06/2015	12.5	11.6	8.8	6.6	11.2	2.2	1.3	1.5	4.0	3.4
16/06/2015	7.3	6.3	2.8	2.7	8.7	5.5	4.6	2.1	7.8	6.5
17/06/2015	7.2	5.7	5.5	1.6	10.2	1.6	-	1.6	3.0	4.4
18/06/2015	6.2	4.4	2.6	2.4	8.1	2.0	-0.1	1.7	3.6	1.1
19/06/2015	6.2	5.4	3.8	1.7	6.6	3.2	0.4	2.3	2.4	2.6
20/06/2015	11.0	9.3	7.5	4.8	7.1	8.0	0.7	1.1	2.5	4.9
21/06/2015	13.7	7.6	9.4	4.0	8.4	2.0	0.3	1.9	-	2.8
22/06/2015	17.2	6.9	15.1	3.2	9.4	0.6	2.7	3.0	3.6	5.2
23/06/2015	20.4	8.4	14.4	3.8	10.0	3.5	2.8	1.2	3.2	7.7
24/06/2015	20.9	9.7	11.7	3.6	12.0	2.1	6.2	3.6	4.3	5.5
25/06/2015	13.4	11.1	7.0	3.9	10.0	4.3	0.5	0.7	1.6	4.7
26/06/2015	15.8	13.3	9.3	4.9	10.0	0.9	1.1	1.6	1.2	1.5
27/06/2015	19.4	9.8	17.2	4.5	9.4	7.0	6.1	1.6	5.5	2.6
28/06/2015	18.2	11.4	17.2	7.8	10.1	1.1	1.4	1.3	1.9	-
29/06/2015	18.8	12.0	13.7	6.5	11.9	1.5	0.7	1.1	0.9	-
30/06/2015	15.7	9.3	12.4	5.6	11.9	3.0	2.0	2.0	4.3	6.3

⁻ Not applicable

Table C-2: June 24-hour average HVAS monitoring data

Table C-2: June 24-hour average HVAS monitoring data							
Date		(HVAS) /m³)					
	Wakefield (Westside)	Teralba					
4/06/2015	9.3	13.0					
10/06/2015	8.5	8.0					
16/06/2015	6.8	8.0					
22/06/2015	10.2	8.0					
28/06/2015	13.9	20.0					

⁻ Not applicable